

American Board of Aesthetic Medicine Board Certification

Dermal Filler Injection: a case report

Performed by: _____, MD

Case Report:

This is a case of a healthy 42 year-old male, Fitzpatrick III, who received injections of HA (Restylane-L (Restylane with lidocaine)) in the midface and nasolabial area with no adverse reaction. Patient had no contraindications, which includes: coagulation disorder, anti-coagulant medication, previous allergic reaction, active herpes simplex infection, or anti-platelet medication or herbal agents. A week later, the nasolabial folds smoothed, with fine line reduction, and improved facial contouring. No adverse reaction at time of application. In image 1a and 1b, the patient has a well-defined volume loss, with intact skin and no signs of infection. In Image 1c and 1d, the patient had reduction of the volume loss, with improvement of facial contouring, and no significant swelling or nodularity. Moreover, the HA injection was placed in the mid-dermis, using a 29 gauge needle, using an overlapping technique, by 1-2mm.

Image 1: a. Pre-operative HA midface injection sites. b. Pre-operative HA right anterolateral view of midface injection sites c. Post-operative HA injections of right and left midface, at 7 days. d. Post-operative HA right anterolateral view of midface injection site, at 7 days.

View: Pre-Operative Post-Operative

Frontal

- a.
- c.

Anterolateral-Right

- b.